

$\nu_e \rightarrow \nu_s$ oscillations with large neutrino mass in NuTeV?

Carlo Giunti

*INFN, Sezione di Torino,**and**Dipartimento di Fisica Teorica, Università di Torino,**Via P. Giuria 1, I-10125 Torino, Italy*

Marco Laveder

*Dipartimento di Fisica “G. Galilei”, Università di Padova,**and**INFN, Sezione di Padova,**Via F. Marzolo 8, I-35131 Padova, Italy***Abstract**

We propose an explanation of the NuTeV anomaly in terms of oscillations of electron neutrinos into sterile neutrinos. We derive an average transition probability $P_{\nu_e \rightarrow \nu_s} \simeq 0.21 \pm 0.07$, which is compatible with other neutrino data if the mass-squared difference that drives the oscillations is $\Delta m^2 \sim 10 - 100 \text{ eV}^2$.

The NuTeV collaboration measured recently [1] a value of the electroweak parameter $\sin^2 \theta_W$ higher than the standard model prediction obtained from a fit of other electroweak data by

$$\sin^2 \theta_W^{\text{NuTeV}} - \sin^2 \theta_W = (5.0 \pm 1.6) \times 10^{-3}. \quad (1)$$

In this paper we propose an explanation of this discrepancy in terms of $\nu_e \rightarrow \nu_s$ oscillations¹, where ν_s is a sterile neutrino that does not participate to weak interactions (see [3, 4]). Our analysis is approximate and based on the limited information contained in the paper of the NuTeV collaboration [1]. A precise analysis of NuTeV data in terms of neutrino oscillations can be done only by the NuTeV collaboration.

The NuTeV collaboration measured the ratio of neutral current to charged current cross sections for neutrino scattering with isoscalar targets of u and d quarks

$$R^\nu = \frac{\sigma(\nu_\mu + N \rightarrow \nu + X)}{\sigma(\nu_\mu + N \rightarrow \mu^- + X)}, \quad (2)$$

and the corresponding ratio $R^{\bar{\nu}}$ for antineutrinos. The Paschos-Wolfenstein relation [5] allows to derive the value of $\sin^2 \theta_W$ from R^ν and $R^{\bar{\nu}}$:

$$\sin^2 \theta_W = \frac{1}{2} - \frac{R^\nu - r R^{\bar{\nu}}}{1 - r}, \quad (3)$$

¹ Other possible explanations of the NuTeV anomaly have been discussed in Ref. [2].

where

$$r \equiv \frac{\sigma(\bar{\nu}_\mu + N \rightarrow \mu^+ + X)}{\sigma(\nu_\mu + N \rightarrow \mu^- + X)} \simeq \frac{1}{2}. \quad (4)$$

The NuTeV collaboration distinguishes CC and NC ν_μ interactions on the basis of the length of the track in the detector: long tracks correspond to CC ν_μ interactions and short tracks correspond to NC ν_μ interactions, plus a contribution from CC and NC ν_e interactions. The number $N_{\nu_e}^{\text{MC}}$ of ν_e interactions calculated through a Monte Carlo is subtracted to the number N_S^ν of short tracks in order to obtain the number of NC ν_μ interactions used to evaluate R^ν :

$$R_{\text{NuTeV}}^\nu = \frac{N_S^\nu - N_{\nu_e}^{\text{MC}}}{N_L^\nu}, \quad (5)$$

where $N_L^\nu = N_{\nu_\mu}^{\text{CC}}$ is the number of long tracks. If $\nu_e \rightarrow \nu_s$ oscillations deplete the number of ν_e interactions with respect to the calculated one, the number $N_{\nu_e}^{\text{MC}}$ subtracted in the numerator of Eq. (5) is too high, leading to a smaller value of R_{NuTeV}^ν with respect to the true one and, from the Paschos-Wolfenstein relation (3), a value of $\sin^2 \theta_W^{\text{NuTeV}}$ larger than the true $\sin^2 \theta_W$.

In the presence of $\nu_e \rightarrow \nu_s$ transitions, the true number of ν_e interactions is

$$N_{\nu_e} = N_{\nu_e}^{\text{MC}} P_{\nu_e \rightarrow \nu_e}, \quad (6)$$

where

$$P_{\nu_e \rightarrow \nu_e} = 1 - P_{\nu_e \rightarrow \nu_s}. \quad (7)$$

is the electron neutrino survival probability averaged over the energy spectrum of the NuTeV beam. Writing

$$N_S^\nu = N_{\nu_\mu}^{\text{NC}} + N_{\nu_e}, \quad (8)$$

from Eq. (5) the ratio R_{NuTeV}^ν reported by the NuTeV collaboration is given by

$$R_{\text{NuTeV}}^\nu = \frac{N_{\nu_\mu}^{\text{NC}}}{N_{\nu_\mu}^{\text{CC}}} - \frac{N_{\nu_e}^{\text{MC}}}{N_{\nu_\mu}^{\text{CC}}} (1 - P_{\nu_e \rightarrow \nu_e}) = R^\nu - \frac{N_{\nu_e}^{\text{MC}}}{N_{\nu_\mu}^{\text{CC}}} (1 - P_{\nu_e \rightarrow \nu_e}), \quad (9)$$

where R^ν is the true value of the ratio (2). A similar relation holds for the antineutrino ratio:

$$R_{\text{NuTeV}}^{\bar{\nu}} = R^{\bar{\nu}} - \frac{N_{\bar{\nu}_e}^{\text{MC}}}{N_{\bar{\nu}_\mu}^{\text{CC}}} (1 - P_{\nu_e \rightarrow \nu_e}), \quad (10)$$

with the same survival probability, because of CPT invariance (we assume that ν_e and $\bar{\nu}_e$ have approximately the same energy spectrum).

The NuTeV collaboration reported that [1]

$$\frac{N_{\nu_e}^{\text{MC}}}{N_L^\nu + N_S^\nu} \simeq \frac{N_{\bar{\nu}_e}^{\text{MC}}}{N_L^{\bar{\nu}} + N_S^{\bar{\nu}}} \simeq 1.7 \times 10^{-2}. \quad (11)$$

In a first approximation we have

$$\frac{N_{\nu_e}^{\text{MC}}}{N_{\nu_\mu}^{\text{CC}}} = \frac{N_{\nu_e}^{\text{MC}}}{N_{\nu_\mu}^{\text{NC}} + N_{\nu_\mu}^{\text{CC}}} \frac{N_{\nu_\mu}^{\text{NC}} + N_{\nu_\mu}^{\text{CC}}}{N_{\nu_\mu}^{\text{CC}}} \simeq \frac{N_{\nu_e}^{\text{MC}}}{N_L^\nu + N_S^\nu} (1 + R_{\text{NuTeV}}^\nu), \quad (12)$$

and a similar relation for $N_{\bar{\nu}_e}^{\text{MC}}/N_{\bar{\nu}_\mu}^{\text{CC}}$. From Eq. (11) and the values $R_{\text{NuTeV}}^\nu \simeq R_{\text{NuTeV}}^{\bar{\nu}} \simeq 0.4$ reported by the NuTeV collaboration [1], we obtain

$$\frac{N_{\nu_e}^{\text{MC}}}{N_{\nu_\mu}^{\text{CC}}} \simeq \frac{N_{\bar{\nu}_e}^{\text{MC}}}{N_{\bar{\nu}_\mu}^{\text{CC}}} \simeq 2.4 \times 10^{-2}. \quad (13)$$

From Eqs. (7), (9), (10), (13) and the Paschos-Wolfenstein relation (3), the value $\sin^2 \theta_W^{\text{NuTeV}}$ reported by the NuTeV collaboration is given by

$$\sin^2 \theta_W^{\text{NuTeV}} = \frac{1}{2} - \frac{R_{\text{NuTeV}}^\nu - r R_{\text{NuTeV}}^{\bar{\nu}}}{1 - r} = \sin^2 \theta_W + \frac{N_{\nu_e}^{\text{MC}}}{N_{\nu_\mu}^{\text{CC}}} P_{\nu_e \rightarrow \nu_s}. \quad (14)$$

Using the numerical values in Eqs. (1) and (13), we finally obtain

$$P_{\nu_e \rightarrow \nu_s} \simeq 0.21 \pm 0.07. \quad (15)$$

The relatively large transition probability (15) is compatible with the fact that reactor experiments (see Ref. [6]) did not measure a disappearance of electron antineutrinos if the mass-squared difference that drives the NuTeV oscillations is²

$$\Delta m^2 \sim 10 - 100 \text{ eV}^2, \quad (16)$$

corresponding to an oscillation length in reactor experiments

$$L_{\text{osc}}^{\text{reactors}} = \frac{4\pi E_{\text{reactors}}}{\Delta m^2} \sim 1 - 10 \text{ cm}, \quad (17)$$

where $E_{\text{reactors}} \sim 1 \text{ MeV}$. Since the source-detector distance in reactor neutrino oscillation experiments is larger than about 10 m, the average survival probability of electron antineutrinos is a constant and cannot be measured without assuming a precise knowledge of the initial $\bar{\nu}_e$ flux.

The oscillation length corresponding to a Δm^2 in the range (16) for an energy of 100 GeV, typical of the NuTeV experiment, is

$$L_{\text{osc}}^{\text{NuTeV}} \sim 1 - 10 \text{ km}, \quad (18)$$

which is appropriate for the observation of transitions with the NuTeV source-detector distance of 1.5 km.

Disappearance of electron neutrinos driven by the Δm^2 given in Eq. (16) is particularly attractive for measurements in the near future using the novel beta beam technique proposed in Ref. [8], with a source-detector distance of the order of 10 m for a neutrino energy $E \gtrsim 0.1 - 1 \text{ GeV}$.

These $\nu_e \rightarrow \nu_s$ transitions could also be observed in other high precision accelerator short-baseline neutrino oscillation experiments like NOMAD [9], with a source-detector distance of the order of 1 km for a neutrino energy $E \gtrsim 10 - 100 \text{ GeV}$.

² Our results may be compatible with those presented a long time ago in Ref. [7].

A high precision measurement could be done in the long term at a neutrino factory (see Ref. [10, 11]) with a short baseline detector located at a source-detector distance of 100 m with a neutrino energy $E \gtrsim 1 - 10 \text{ GeV}$ [12, 13].

The Δm^2 in Eq. (16) could be the same that drives the small $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations observed in the LSND experiment [14]. In this case, $\nu_\mu \rightarrow \nu_e$ appearance will be observed in the coming MiniBooNE experiment [15]. Simultaneous $\nu_e \rightarrow \nu_s$ disappearance and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance from neutrinos produced in $\mu^+ \rightarrow \bar{\nu}_\mu + e^+ + \nu_e$ decays³ driven by the same Δm^2 given in Eq. (16) could be easily observed at a neutrino factory with a short baseline detector with electron charge discrimination capability [12]. The signature of $\nu_e \rightarrow \nu_s$ oscillations will be a depletion in the e^- energy spectrum induced by ν_e 's and the presence of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations will be characterized unequivocally by the appearance of e^+ events.

$\nu_e \rightarrow \nu_s$ transitions driven by the high Δm^2 in Eq. (16) contribute with an energy-independent probability to the disappearance of solar electron neutrinos and must be taken into account in the analysis of solar neutrino data (see [3, 4]). The energy dependence of the solar electron neutrino disappearance probability observed in solar neutrino experiments could be due to another very small squared-mass difference, in the framework of four-neutrino mixing schemes compatible also with the experimental evidence of atmospheric neutrino oscillations (see [3, 4, 16]).

The range (16) for Δm^2 is compatible with the upper limit $m_{\nu_e} \lesssim 3 \text{ eV}$ found in tritium decay experiments (see Ref. [17]). Indeed, the averaged probability (15) implies approximately

$$\sin^2 2\vartheta \sim 0.4, \quad (19)$$

where ϑ is the mixing angle. Assuming a hierarchy of neutrino masses, the effective electron neutrino mass in tritium beta-decay experiments is

$$m_{\nu_e} \simeq \sin^2 \vartheta \sqrt{\Delta m^2} \sim 0.3 - 1 \text{ eV}, \quad (20)$$

that is compatible with the experimental upper bound.

If there is a hierarchy of neutrino masses, the effective neutrino mass in neutrinoless double-beta decay is equal to m_{ν_e} in Eq. (20). This value is compatible with the existing experimental upper limits [18, 19] and curiously overlaps with the evidence in favor of neutrinoless double-beta decay claimed in Ref. [20] (see, however, the discussion in Refs. [21, 22]).

The large neutrino mass implied by the Δm^2 in Eq. (16) is compatible with the cosmological upper bound on light neutrino masses (about 10 eV), but would have important effects on the evolution of the universe (see Ref. [23]).

In conclusion, we have proposed an explanation of the NuTeV in terms of $\nu_e \rightarrow \nu_s$ oscillations, with the relatively large average transition probability given in Eq. (15) and a large squared-mass difference given in Eq. (16). These transitions are compatible with the results of other neutrino experiments and could be verified by future high-precision experiments. Let us finally remark again that our analysis has been approximate and based on the limited information available in the paper of the NuTeV collaboration [1]. We hope that the NuTeV collaboration will consider seriously our suggestion and perform a precise analysis of their data, which could lead to a determination of the allowed ranges of Δm^2 and $\sin^2 2\vartheta$.

³ Or $\bar{\nu}_e \rightarrow \bar{\nu}_s$ disappearance and $\nu_\mu \rightarrow \nu_e$ appearance from neutrinos produced in $\mu^- \rightarrow \nu_\mu + e^- + \bar{\nu}_e$ decays.

References

- [1] NuTeV, G. P. Zeller *et al.*, (2001), hep-ex/0110059.
- [2] S. Davidson, S. Forte, P. Gambino, N. Rius, and A. Strumia, (2001), hep-ph/0112302.
- [3] S. M. Bilenky, C. Giunti, and W. Grimus, Prog. Part. Nucl. Phys. **43**, 1 (1999), hep-ph/9812360.
- [4] M. C. Gonzalez-Garcia and Y. Nir, (2002), hep-ph/0202058.
- [5] E. A. Paschos and L. Wolfenstein, Phys. Rev. **D7**, 91 (1973).
- [6] C. Bemporad, G. Gratta, and P. Vogel, (2001), hep-ph/0107277.
- [7] G. Conforto, Nuovo Cim. **A103**, 751 (1990).
- [8] P. Zucchelli, (2001), hep-ex/0107006.
- [9] NOMAD, P. Astier *et al.*, Nucl. Phys. **B611**, 3 (2001), hep-ex/0106102.
- [10] C. Albright *et al.*, (2000), arXiv:hep-ex/0008064.
- [11] A. Blondel *et al.*, Nucl. Instrum. Meth. **A451**, 102 (2000).
- [12] A. Bueno, M. Campanelli, M. Laveder, J. Rico, and A. Rubbia, JHEP **06**, 032 (2001), hep-ph/0010308.
- [13] T. Adams *et al.*, (2001), hep-ph/0111030.
- [14] LSND, A. Aguilar *et al.*, Phys. Rev. **D64**, 112007 (2001), hep-ex/0104049.
- [15] BooNe, E. Church *et al.*, FERMILAB-P-0898.
- [16] C. Giunti, M. Laveder, and M. Mezzetto, (2002), Neutrino Unbound WWW page: <http://www.to.infn.it/~giunti/NU>.
- [17] D. E. Groom *et al.*, Eur. Phys. J. **C15**, 1 (2000), WWW page: <http://pdg.lbl.gov>.
- [18] H. V. Klapdor-Kleingrothaus *et al.*, Eur. Phys. J. **A12**, 147 (2001).
- [19] C. E. Aalseth *et al.*, (2002), hep-ex/0202026.
- [20] H. V. Klapdor-Kleingrothaus, A. Dietz, H. L. Harney, and I. V. Krivosheina, Mod. Phys. Lett. **A16**, 2409 (2002), hep-ph/0201231.
- [21] C. E. Aalseth *et al.*, (2002), hep-ex/0202018.
- [22] F. Feruglio, A. Strumia, and F. Vissani, (2002), hep-ph/0201291.
- [23] A. D. Dolgov, (2002), hep-ph/0202122.